

TRANSFER REACTIONS

THE STRUCTURE OF ^{18}Ne AND THE BREAKOUT OF THE HOT CNO CYCLE

K.I. Hahn and N. Bateman

Yale University, New Haven, Connecticut 06511

P. Magnus

University of Washington, Seattle, Washington 98195

A. Bacher, G.P.A. Berg, and E.J. Stephenson

Indiana University Cyclotron Facility, Bloomington, Indiana 47408

The nuclear structure of ^{18}Ne plays an important role in explosive hydrogen burning events. Both the $^{14}\text{O}(\alpha, p)^{17}\text{F}$ and $^{17}\text{F}(p, \gamma)^{18}\text{Ne}$ reactions proceed predominantly through resonances in ^{18}Ne . $^{14}\text{O}(\alpha, p)^{17}\text{F}$ plays an important role in the breakout from the astrophysical hot CNO cycle and $^{17}\text{F}(p, \gamma)^{18}\text{Ne}$ plays the critical role in determining the ratio of $A=17$ to $A=18$ nuclides produced in explosive hydrogen burning. Ideally these rates would be determined by measuring $^{14}\text{O}+\alpha$ and $^{19}\text{F}+p$ directly, but that would require an enormous effort since $^{14}\text{O}(t_{1/2}=71\text{ s})$ and $^{17}\text{F}(t_{1/2}=65\text{ s})$ are unstable. Another alternative is to determine the rate by measuring the appropriate resonance parameters in ^{18}Ne . There are two light ion reactions with which the structure of ^{18}Ne can be studied with high resolution, $^{20}\text{Ne}(p, t)$ and $^{16}\text{O}(^3\text{He}, t)$. We studied the excitation range of interest using the $^{16}\text{O}(^3\text{He}, t)$ reaction at the University of Washington and the $^{20}\text{Ne}(p, t)$ reaction at IUCF using the K600 spectrometer. We report on the latter here.

We measured the $^{20}\text{Ne}(p, t)^{18}\text{Ne}$ reaction at 6° and 11° using a 90 MeV proton beam and the K600 spectrometer in order to determine the excitation energies and, where possible, the natural widths of levels between 6 and 8 MeV of excitation, and in an attempt to confirm the observation¹ of the $J^\pi = 3^+$ level at 4.56 MeV. Figure 1 shows the spectrum taken at 6° . Critical to the measurement was the target made by implanting $7\text{ }\mu\text{g}/\text{cm}^2$ ^{20}Ne into natural $40\text{ }\mu\text{g}/\text{cm}^2$ carbon foils.² This enabled us to have a 25-30 keV resolution which would not have been possible with a standard gas cell.

The analysis of these data is not complete, but preliminary results show two interesting points. First we saw only two levels in the 6.0-6.5 MeV excitation energy range, the range critical to the $^{14}\text{O}+\alpha$ reaction (Fig. 1), at 6.30 and 6.35 MeV of excitation. These two levels and one other at 6.15 MeV were seen in this excitation range in $^{16}\text{O}(^3\text{He}, n)^{18}\text{Ne}$ (see Ref. 3) and also in a heavy-ion reaction of $^{12}\text{C}(^{12}\text{C}, ^6\text{He})^{18}\text{Ne}$ (see Ref. 4). There are only three levels in the well-studied analog nucleus ^{18}O in this excitation range. They have $J^\pi=1^-, 2^-$ and 3^- . Considering that the $^{20}\text{Ne}(p, t)$ reaction strongly prefers to populate natural parity states, we tentatively conclude that the 6.15 MeV level in ^{18}Ne not seen in




Figure 1. The $^{20}\text{Ne}(p,t)^{18}\text{Ne}$ spectrum taken at a beam energy of 90 MeV and an angle of 6° . The peaks resulting from $^{20}\text{Ne}(p,t)^{18}\text{Ne}$ are labeled by the excitation of the state in ^{18}Ne populated. Peaks that result from $^{12}\text{C}(p,t)^{10}\text{C}$ and $^{13}\text{C}(p,t)^{11}\text{C}$ in the target are labeled with ^{10}C and ^{11}C respectively.

(p,t) is the analog of the 2^- in ^{18}O . This is in disagreement with the conclusions of Garcia who tentatively assigns $J^\pi=1^-$ to this level based on Coulomb shift calculations. Since Garcia calculates that the 1^- level dominates the $^{14}\text{O}+\alpha$ reaction rate, these (p,t) results require a recalculation of that reaction rate which is now in progress.

Secondly we do not see any evidence for the 3^+ level at 4.56 MeV, which is not surprising considering the strong preference the (p,t) reaction has for populating natural parity levels.

1. A. Garcia *et al.*, Phys. Rev. **C43**, 2012 (1991).
2. M. Smith *et al.*, Nucl. Phys. **A536**, 333 (1992).
3. A. Garcia, Thesis, University of Washington 1991.
4. K.I. Hahn, Thesis, Yale University (in progress) 1992.